

Resource for Management Options of Endocrine Cancer During COVID-19

DATE: March 23, 2020



SSO supports the need for treatment decisions to be made on a case-by-case basis. The surgeon's knowledge and understanding of the biology of each cancer, alternative treatment options, and the institution's policies at the time the patient will be scheduled for surgery all need to be taken into consideration.

The information below is based on the opinions of individuals who are experts within the field of endocrine cancer and are members of the Society's Endocrine/Head and Neck Disease Site Work Group.

The opinions generally align or incorporate those from other societies such as the American Association of Endocrine Surgeons (AAES).

- Most uncomplicated endocrine operations can be delayed. Diseases and presentations that might qualify for more **urgent surgery** (i.e., within approximately 4-8 weeks during the current pandemic), include:
 - Thyroid:
 - Thyroid cancer that is a current or impending threat to life, those that are threatening morbidity with local invasion (e.g., trachea, recurrent laryngeal nerve), aggressive biology (rapidly growing tumor or recurrence, rapidly progressive local-regional disease including lymph nodes)
 - Severely symptomatic Graves' disease that has failed medical therapy
 - Goiter that is highly symptomatic or is at risk for impending airway obstruction
 - Open biopsy with diagnostic intent for suspected anaplastic thyroid cancer or lymphoma
 - Parathyroid:
 - Hyperparathyroidism with life-threatening hypercalcemia that cannot be controlled medically
 - Adrenal:
 - Adrenocortical cancer or highly suspected adrenocortical cancer
 - Pheochromocytoma or paraganglioma that is unable to be controlled with medical management
 - Cushing's syndrome with significant symptoms that is unable to be controlled with medical management
 - Generally, functional adrenal tumors that are medically controlled and asymptomatic non-functional adrenal adenomas can be delayed
 - Neuroendocrine Tumors:
 - Symptomatic small bowel NETs (e.g., obstruction, bleeding/hemorrhage, significant pain, concern for ischemia)
 - Symptomatic and/or functional pancreatic NETs that cannot be controlled medically
 - Non-functional pancreatic NETs causing symptoms (jaundice, bleeding, obstruction) after failure of somatostatin analogues and medical therapy

- Lesions with significant growth or short doubling times
- Most other NETs can be delayed and/or can be considered for somatostatin analogues or medical therapy
- Cytoreductive operations and metastasectomy should generally be delayed but should be considered on an individual basis, especially if progressing after multiple therapies (SSAs, chemo or biologic therapy, PRRT)

Other:

- If an endocrine disorder threatens a pregnant mother or her fetus

A. Surveillance for delayed procedures:

Adrenal:

- CT or MRI every 3-6 months
- Functional tumors:
 - Appropriate laboratory studies every 3-6 months
 - Medical management of functional tumors should be ongoing

Parathyroid:

- Medical management when necessary

Thyroid:

- Ultrasound surveillance (including nodal basins) and physical exam every 3-6 months

Neuroendocrine Tumors:

- CT or MRI imaging every 3-6 months
- Functional tumors:
 - Appropriate laboratory studies every 3-6 months
 - Medical management of functional tumors should be ongoing

B. Endoscopic Surveillance for gastric, duodenal, or rectal NET

- Consider deferring as the emerging evidence on the risk of transmission of SARS-CoV-2 with procedures involving the GI tract should be considered for the safety of both the surgical team and the patient.

Safety Considerations for Surgeons Performing Head and Neck Procedures:

- The evidence suggesting that surgeons performing procedures on the head and neck are at higher risk for COVID-19 should be reviewed and considered.
- Appropriate PPE for any surgical procedure should always be utilized and should conform to the recommendations of the CDC and your local institution.
- Avoid or minimize exposure to any Aerosol-Generating procedures such as intubation/extubation, as these are high risk situations for SARS-CoV-2 transmission.